

GRAVITATION SINGULARITIES OF THE CAUSTIC TYPE

G.Sardanashvily

Department of Theoretical Physics, Moscow State University
117234 Moscow, Russia

Various criterion of gravitational singularities have been suggested [3, 6, 7]. In view of the well-known correspondence between the space-time distributions and the gravitational fields on a world manifold X^4 , gravitational singularities can be indicated by singularities of these distributions [10, 11]. In the germ terms, singularities of a (3+1) distribution look locally like singularities of a foliation whose leaves are level surfaces of a real function f on X . If f is a single-valued function, changes of leave topology at critical points of f take place [10]. In case of a multi-valued function f , one can lift the foliation to the total space of the cotangent t^*X bundle over X , then extend it over branch points of f and project this extension onto X . Singular points of this projection constitute a Lagrange map caustic by Arnol'd.

In gravitational theory, a space-time structure is usually defined to be a (3+1) decomposition

$$TX = \mathbf{F} \oplus T^0X \quad (1)$$

of the tangent bundle over a world manifold into a 3-dimensional spatial sub-bundle $\mathbf{F}$ and its time-like orthocomplement T^0X .

Recall that there is the 1:1 correspondence between the nonvanishing 1-forms ω on a manifold X and the smooth orientable distributions $\mathbf{F}$ of 1-codimensional subspaces of tangent spaces to X . This correspondence is defined by the equation

$$\mathbf{F} \rfloor \omega = 0.$$

A form ω is called a generating form of a distribution $\mathbf{F}$.

We have the following theorem [7, 12] formulated in terms of space-time distributions [11].

THEOREM F. *For every gravitational field g on a world manifold X , there exists an associated pair $(\mathbf{F}, g^R)$ of a space-time distribution $\mathbf{F}$ with a generating*

tetrad form

$$h^0 = h_\mu^0 dx^\mu$$

and a Riemannian metric g , so that

$$g^R = 2h^0 \otimes h^0 - g = h^0 \otimes h^0 + k \quad (2)$$

where k is a Riemannian metric in the tangent subbundle $\mathbf{F}$. Conversely, given a Riemannian metric g^R , every oriented smooth 3-dimensional distribution $\mathbf{F}$ with a generating form ω is a space-time distribution compatible with the gravitational field g given by expression (2) where

$$h^0 = \omega / |\omega|, \quad |\omega|^2 = g^R(\omega, \omega) = g(\omega, \omega).$$

A 1-codimensional distribution $\mathbf{F}$ is called an integrable distribution if its generating form ω obeys the equation

$$\omega \wedge d\omega = 0.$$

In this case, fibres of the corresponding (3+1) decomposition (1) are tangent to leaves of some 1-codimensional foliation of spatial hypersurfaces of a world manifold X . An integrable space-time distribution (a space-time foliation) is called causal if its generating form can be exact, that is,

$$\omega = df$$

where f is some real function on X which has no critical points where $df = 0$ [8]. This notion of causality coincides with the one of stable causality by Hawking [7]. Leaves of a causal foliation are level surfaces of its generating function f . No curve transversal to leaves of a causal foliation intersects each leave more than once.

We say that a gravitational field g on X is free from singularities if there exists an associated pair of a complete Riemannian metric g^R and a causal space-time foliation $\mathbf{F}$ with the generating form h^0 such that $g^R(\nabla h^0, \nabla h^0)$ is bounded on X . This condition guaranties that, being complete with respect to g^R , a space-time satisfies the well-known b -completeness condition [2].

One can distinguish several types of gravitation singularities in accordance to this criterion. We examine gravitation singularities characterized by singularities of space-time distributions. The distribution singularities can be

described locally (in the germ form) as singularities of a causal foliation with a generating function f . There are two types of these singularities.

(i) A single-valued generating function f has critical points where $df = 0$. It generates the Haefliger structure (the singular foliation) of its level sets on X . These level sets change their topology at critical points of f . Gravitation singularities of this type are the scalar curvature singularities in accordance to the classification in [4].

(ii) A generating function f is a multiple-valued function on X . The leaves of the foliation $\mathbf{F}$ defined on the domain where f is a single-valued begin to intersect each other at branch points of f . Branch points of f where the foliation is destroyed form a caustic. To describe foliation singularities of this type, one can lift the space-time foliation $\mathbf{F}$ into the total space of the cotangent bundle T^*X , then extend this lifted foliation over the singular points, and project the extended foliation onto the base X . Singularities of $\mathbf{F}$ can be described as singularities of this projection.

In gravitation theory, a geometrical locus of focal and conjugate points is called a caustic by analogy with geometrical optics [9, 13]. We follow the general mathematical notion of caustics as singularities of the Lagrange maps [1, 5]. Each caustic can be brought locally (in germ terms) into the following standard form.

Let a space $\mathbf{R}^{2n}$ be endowed with the coordinates $\{x^\mu, P_\mu\}$. Let us consider the Liouville form

$$\alpha = P_\mu dx^\mu \quad (3)$$

on $\mathbf{R}^{2n}$ and a submanifold N of $\mathbf{R}^{2n}$ such that

$$d\alpha(N) = 0,$$

that is, being restricted onto N , the form α is exact:

$$\alpha(N) = dz(N).$$

Such a manifold of maximal dimension n is called a Lagrange submanifold. A Lagrange submanifold can be defined by a generating function $S(x^i, P_j)$ of n variables $(x^i, P_j, i \in I, j \in J)$ (where (I, J) is some partition of the set $(1, \dots, n)$). It is given by the relations

$$x^j = -\frac{\partial S}{\partial P_j}, \quad P_i = \frac{\partial S}{\partial x^i}.$$

Let us consider the projection

$$\pi : (x^\mu, P_\mu) \rightarrow (x^\mu)$$

of $\mathbf{R}^{2n}$ onto $\mathbf{R}^n$. Being restricted to the Lagrange submanifold

$$\pi_N : (x^i, P_j) \rightarrow (x^i, x^j = -\frac{\partial S}{\partial P_j}),$$

this projection is called the Lagrange map. A caustic is defined to be the set of critical points of a Lagrange map, i.e., the points where the matrix

$$\partial^2 S / \partial P_i \partial P_j$$

is singular.

For instance, a caustic on manifolds is defined as follows. Let the cotangent bundle T^*X be provided with the induced coordinates $(x^\mu, P_\mu, \dots x_\mu)$. The Liouville form (3) defines n -dimensional Lagrange submanifolds of T^*X . Singular points of projection of such a Lagrange submanifold onto the base X form a caustic.

Let us note that a geometrical locus of focal and conjugate points of Riemannian and time-like pseudo-Riemannian geodesics also is a caustic in accordance to the Arnol'd definition [10].

Our definition of foliation caustics is based on the following proposition.

PROPOSITION . *For any foliation of level surfaces $\mathbf{F}$ of a manifold X , there is a foliation $\mathbf{F}'$ of some Lagrange submanifold of T^*X such that $\mathbf{F}$ is the image of $\mathbf{F}'$ under the Lagrange map.*

Proof. Let f be a generating function of the foliation $\mathbf{F}$. We define the embedding

$$\gamma : (x^\mu) \rightarrow (x^\mu, P_\mu = \frac{\partial f}{\partial x^\mu})$$

of X into T^*X . Its image is a Lagrange submanifold of $T^*(X)$. Let $\mathbf{F}'$ be the induced foliation $\pi_{\gamma(X)}^* \mathbf{F}$ of $\gamma(X)$ where $\pi_{\gamma(X)}$ is the Lagrange map

$$\pi_{\gamma(X)} : \gamma(X) \rightarrow X.$$

Since γ and $\pi_{\gamma(X)}$ are diffeomorphisms between X and $\gamma(X)$ ($\pi_{\gamma(X)} \circ \gamma = \text{id}_X$), the foliation $\mathbf{F}$ on X can be represented as the image of the foliation $\mathbf{F}'$ on $\gamma(X)$ under the Lagrange map $\pi_{\gamma(X)}$.

For instance, let $N \subset T^*X$ be the Lagrange submanifold generated locally by a function $S(x^i, P_j)$, and let $\mathbf{F}'$ be the foliation of level surfaces of the function

$$f'(x^i, P_j) = S - P_j \frac{\partial S}{\partial P_j}$$

on the Lagrange submanifold N . The image $\pi_N(\mathbf{F}')$ of $\mathbf{F}'$ under the Lagrange map π_N is a foliation of the image $\pi_N(U)$ of the domain $U \subset N$ where this Lagrange map has no critical points. This foliation is destroyed at caustic points of the Lagrange map π_N .

There are the following classes $A_2, A_3, A_4, D_4, A_5, D_5$ of stable caustics on a 4 dimensional manifold. For example, the canonical generating function of caustic A_3 takes the form

$$S = -P_0^4 + x^1 P_0^2,$$

and the corresponding Lagrange manifold N is given by equations

$$x^0 = 4P_0^3 - 2x^1 P_0, \quad P_1 = P_0^2.$$

The Lagrange map then reads

$$x^0 = 4P_0^3 - 2x^1 P_0. \tag{4}$$

The caustic set where

$$\partial^2 S / \partial P_0^2 = 0$$

consists of the points

$$x^1 = 6P_0^2,$$

and its Lagrange image on X contains the points

$$(x^0)^2 = \frac{8}{27}(x^1)^5.$$

The generating function of the foliation $\mathbf{F}'$ on the Lagrange manifold takes the form

$$f' = 3P_0^4 - x^1 P_0^2.$$

Then, on X , the generating function of the Lagrange image of $\mathbf{F}'$ reads

$$f(x^0, x^1) = f'(x^1, P_0(x^0, x^1))$$

where the function $P_0(x^0, x^1)$ is defined by equation (4). The function P_0 and the generating function f become three-valued functions at caustic points. The A_3 -germ of foliation caustics thus is characterized by the behavior of the component ω_0 of the foliation generating form which is tripled at caustic points.

Caustic singularities have the following feature. There are domains of a space-time where not nearest, but the far separated leaves begin to intersect each other. Therefore, a space-time foliation can be locally prolonged over the caustic points, whereas global prolongation of this foliation is impossible.

For example, let $f(u, v)$ be a real function on $\mathbf{R}^2$ which obeys the equation

$$f^3(u, v) - 3uf(u, v) - 2v = 0$$

where u, v are coordinates on $\mathbf{R}^2$. This function is the singled-valued one

$$f_+ = [v + (v^2 - u^3)^{1/2}]^{1/3} + [v - (v^2 - u^3)^{1/2}]^{1/3}$$

on the domain $U = (u, v : v^2 > u^3)$, and it is the three-valued function

$$\begin{aligned} f_{0,1,2} &= 2u^{1/2} \cos\left(\frac{1}{3}(\phi + 2\pi n)\right), \\ \phi &= \arccos(vu^{-3/2}), \quad n = 0, 1, 2, \end{aligned}$$

at points $v^2 < u^3$. Let $\mathbf{F}$ be the foliation

$$\mathbf{F}_c = \{u, v : f_+(u, v) = c = \text{const}\}$$

on $U \subset \mathbf{R}^2$. Its leaves $\mathbf{F}_c$ are the lines

$$2v = c^3 - 3uc, \quad -\infty < c < +\infty.$$

This foliation has the caustic singularity at the branch points $v^2 = u^3$ of the function f . Moreover, $u = v = 0$ is the A_3 -caustic point, whereas the other ones $v^2 = u^3 \neq 0$ are points of the A_2 -caustic. The leaves $\mathbf{F}_c, c > \alpha > 0$, can be prolonged over the caustic curve $v = u^{3/2}$ onto the domain $0 < v < u^{3/2}$ where they can be described as leaves of the foliation

$$f_0(u, v) = \text{const}.$$

These leaves however begin to intersect each other when $v < 0$, although the nearest leaves intersect each other only on the caustic curve $v = -u^{3/2}$. Note

that the leaves $\mathbf{F}_{c>0}$ begin to intersect the leaves $\mathbf{F}_{c<0}$ on the caustic curve $v = u^{3/2}$.

Caustic singularities however are not reduced to the locally extendible singularities [4]. For instance, the A_2 -caustic points $u^2 = v^3 \neq 0$ of the above-mentioned foliation caustic are locally extendible singularity points, whereas the A_3 -caustic point $u = v = 0$ is not locally extendible.

References

1. V. Arnol'd, *Mathematical Methods in Classical Mechanics* (Springer, Berlin, 1978)
2. M. Canarutto, *Nuovo Cimento*, **80B**, 121, 1984
3. M. Canarutto, *Riv. Nuovo Cimento*, **11**, 1, 1988
4. G. Ellis and B. Schmidt, *GRG*, **10**, 989, 1979
5. H. Friedrich and J. Stewart, *Proc. R. Soc.*, **A385**, 345, 1983..
6. H. Fuchs, U. Kasper, D-E. Liebscher, V. Muller and H-J. Schmidt, *Fortsch. Physik*, **36**, 427, 1988
7. S. Hawking and G. Ellis, *The Large Scale Structure of a Space-Time* (Cambr. Univ. Press, Cambridge, 1973)
8. N. Lawson, *Bull. Amer. Math. Soc.*, **80**, 369, 1974
9. K. Rosquist, *Comm. Math. Phys.*, **88**, 339, 1983
10. G. Sardanashvily and V. Yanchevsky, *Acta Phys. Polon.*, **B17**, 1017, 1986
11. G. Sardanashvily and O. Zakharov, *Gauge Gravitation Theory* (World Scientific, Singapore, 1992).
12. R. Sulanke and P. Wintgen, *Differenialgeometrie und Faserbündel* (Veb Deutscher Verlag der Wissenschaften, Berlin, 1972).
13. F. Warner, *Amer. J. Math.*, **87**, 575, 1965